

Product Information

FSL-1-Cy5


For Research Purposes only. Not for use in Humans



Product	L7050
Chemical name	S-[2,3-bis(palmitoyloxy)-(2RS)-propyl]-(R)-cysteiny-GDPKHPKSFK(Cy5)
Synonyms	Pam ₂ Cys-GDPKHPKSFK(Cyanine5); Cyanine5-labelled Fibroblast-stimulating Lipopeptide
CAS	Not available
MW / Formula	2167 / C ₁₁₆ H ₁₇₇ N ₁₆ O ₁₉ S

Vial content 100 µg

Description



FSL-1-Cy5 is a selectively labelled analogue of FSL-1 (product code L7000). It is labelled with cyanine5 carboxylic acid via the side chain of the C-terminal lysine. Cy5 has an excitation maximum of 646 nm and an emission maximum of 662 nm.

Lipopeptides are valuable tools for basic research in innate and acquired immunity. The synthetic lipopeptide FSL-1 represents the N-terminal sequence of the 44-kDa lipoprotein

LP44 of *Mycoplasma salivarium*. It carries two ester bound fatty acids and a free amino terminus. The synthetic lipopeptide FSL-1 is described to elicit cellular responses through TLR2/TLR6 heterodimers which involves downstream NF-κB activation and cytokine release. The water soluble FSL-1 showed high activity when tested for its capability to activate THP-1 cells to produce TNF-α and on HEK293 cells transfected with TLR2 and TLR6 to produce NF-κB.

Packaging Reconstitution Storage

The lipopeptide is provided as a lyophilised, blue powder without any additives. It can be shipped at room temperature and should be stored at 4°C.

FSL-1-Cy5 can be reconstituted in endotoxin-free water (1 mg/ml stock solution). Through the use of either a homogeniser or sonicator, a homogenous solution or emulsion can be prepared. If you use an ultrasonic bath, take care of the vial labels. For further dilutions water, saline, buffer or media can be used.

After reconstitution, the solution should be aliquoted and stored at or below -20°C. Repeated thawing and freezing should be avoided.

Handling

Good laboratory technique should be employed in the safe handling of any lipopeptide product. If you are not fully trained or are unaware of the hazards involved, do not use this compound!

Caution: Do not take internally! Avoid contact by all modes of exposure. Wear appropriate laboratory attire including a lab coat, gloves, mask and safety glasses. Do not mouth pipette, inhale, ingest or allow coming into contact with open wounds. Wash thoroughly any area of the body which comes into contact with the product. Avoid accidental autoinoculation by exercising extreme care when handling in conjunction with any injection device.

This product is intended for research purposes by qualified personnel only. It is not intended for use in humans or as a diagnostic agent. EMC microcollections GmbH is not liable for any damages resulting from misuse or handling of this product.

Product Information

FSL-1-Cy5

For Research Purposes only. Not for use in Humans



References

- A. O. Aliprantis, R. B. Yang, M. R. Mark, S. Suggett, B. Devaux, J. D. Radolf, G. P. Klimpel, P. Godowski, A. Zychlinsky (1999) Cell Activation and Apoptosis by Bacterial Lipoproteins Through Toll-like Receptor-2. *Science* 285, 736-739.
- K.-I. Shibata, A. Hasebe, T. Into, M. Yamada, T. Watanabe (2000) The N-terminal lipopeptide of a 44 kDa membrane-bound lipoprotein of *Mycoplasma salivarium* is responsible for the expression of intracellular adhesion molecule-1 on the cell surface of normal human gingival fibroblasts. *J. Immunol.* 165, 6538-6544.
- O. Takeuchi, T. Kawai, P. F. Mühlradt, M. Morr, J. D. Radolf, A. Zychlinsky, K. Takeda, S. Akira (2001) Discrimination of bacterial lipoproteins by Toll-like receptor 6. *Int. Immunol.* 13, 933-940.
- O. Takeuchi, S. Sato, T. Horiuchi, K. Hoshino, K. Takeda, Z. Dong, R. L. Modlin, S. Akira (2002) Cutting edge: role of Toll-like receptor 1 in mediating immune response to microbial lipoproteins. *J. Immunol.* 169(1), 10-14.
- J. Nakamura, K. Shibata, A. Hasebe, T. Into, T. Watanabe, N. Ohata (2002) Signaling pathways induced by lipoproteins derived from *Mycoplasma salivarium* and a synthetic lipopeptide (FSL-1) in normal human gingival fibroblasts. *Microbiol. Immunol.* 46, 151-158.
- S. Akira, S. Mammalian (2003) Toll-like receptors. *Curr. Opin. Immunol.* 15, 5-11.
- T. Okusawa, M. Fujita, J.-I. Nakamura, T. Into, M. Yasuda, A. Yoshimura, Y. Hara, A. Hasebe, D. T. Golenbock, M. Morita, Y. Kuroki, T. Ogawa, K.-I. Shibata (2004) Relationship between structures and biological activities of mycoplasmal diacylated lipopeptides and their recognition by Toll-like receptors 2 and 6. *Infect. Immun.* 72(3), 1657-1665.
- K. O. Omueti, J. M. Beyer, C. M. Johnson, E. A. Lyle, R. I. Tapping (2005) Domain exchange between human toll-like receptors 1 and 6 reveals a region required for lipopeptide discrimination. *J. Biol. Chem.* 280, 36616-36625.
- J. Y. Kang, X. Nan, M. S. Jin, S. J. Youn, Y. H. Ryu, S. Mah, S. H. Han, H. Lee, S. G. Paik, J. O. Lee (2009) Recognition of lipopeptide patterns by Toll-like receptor 2-Toll-like receptor 6 heterodimer. *Immunity* doi:10.1016/j.immuni.2009.09.018
- H. Kumar, T. Kawai, S. Akira (2009) Toll-like receptors and innate immunity. *Biochem. Biophys. Res. Comm.* 388, 621-625.
- A. Burger-Kentischer, I. S. Abele, D. Finkelmeier, K.-H. Wiesmüller, S. Rupp (2010) A new cell-based innate immune receptor assay for the examination of receptor activity, ligand specificity, signalling pathways and the detection of pyrogens. *J. Immunol. Methods* 30, 93-103.